

Retreat at Dunsmuir Lodge: A successful and productive methodology workshop

By Amy Poon

On Sunday, April 30, 2000, those from the Fisheries Centre who are involved in the Sea Around Us Project (SAUP) gathered at Dunsmuir Lodge, near Vancouver Island's Swartz Bay, for the welcome dinner (the first of the legendarily divine meals at Dunsmuir) to the SAUP Methodology Workshop. They were joined by six reviewers whom they invited as external auditors to give insights and validity to their work.

The workshop proper, which lasted from Monday, May 1 to Friday, May 5, allowed those involved in the SAUP to get an idea of how their own project fit into the larger scheme of things, as well as get feedback from those who have not worked on the projects on an intimate level. The visitors each brought a different perspective to the proceedings. Lee Alverson (President, Natural Resources Consultants Inc., and Professor of Fisheries, University of Washington) brought with him his extensive experience with

fisheries management. Paul Fanning (Fisheries and Oceans Canada, Halifax) and Poul Degnbol (Director, Institute for Fisheries Management and Coastal Community Development, Denmark) had perspectives from the two sides of the Atlantic. Kevern Cochrane and Richard Grainger (both of the UN Food and Agriculture Organisation in Rome) had insights into international standards, and Jay Maclean (freelance writer and editor) kept the focus on the project's objective. With the added supply of contact names, advice, and suggestions from the visitors, the SAUP's already impressive arsenal was improved.

Because time constraints and lack of personnel were the recurring concerns about the project, some rearrangement of priorities of the SAUP was necessary. Two projects were moved from the SAUP to be picked up later as Fisheries Centre projects. Data analysis would be done on varying levels, where detailed models will come

out of locations in the North Atlantic where data is rich, whereas more general and robust models that could be applied worldwide will come out of locations where data is poor. These, along with other useful suggestions, came out of the visitors' insights.

Although some of the reviewers were initially hesitant about the scope and purpose of the SAUP, they were soon won over. The project was lauded for its holistic and broad approach, its leadership, and its innovation. Concerns such as potential gaps in data, target audience for the reports, and degree of depth, were also addressed.

There is less than a year to go on the SAUP. As the clock ticks, the participants will feel the time crunch more keenly. However, their tasks will be aided by the ideas and suggestions that came out of this workshop.

Amy Poon is an M.Sc. student at the Fisheries Centre, and, along with Yvette Rizzo, acted as rapporteur at the SAUP workshop.

The Sea Around Us Project Newsletter

Issue 4 – May/June 2000

How life history patterns and depth zone analysis can help fisheries policy

By Dirk Zeller and Daniel Pauly

This is a summary of: D. Zeller and D. Pauly (2000) How life history patterns and depth zone analysis can help fisheries policy. In: Pauly, D. and Pitcher, T.J. (eds) Methods for assessing the impact of fisheries on marine ecosystems of the North Atlantic. Fisheries Centre Research Reports 8(1): (in prep).

The stocks of an exploited fish species may be utilized by more than one fishery sector during different stages in the species life history. Often, scientists and managers alike view life history patterns as a multi-dimensional

problem, with complex interactions between components defined by ecology, time and oceanography. Often this complexity has made it difficult to assimilate effects of multiple fishery sectors on a species and the industry it supports. This may be either due to the perception of multi-dimensional complexity thought to be intractable, or because of an oversight of basic patterns.

The life-history patterns of fish species are complex. But much of this complexity can be captured in simple diagrams of coastal transects, where juveniles usually occur inshore in large numbers, while adults are often in deeper, offshore waters. Here we argue that this multi-dimensional complexity can be reduced to a simpler, two-dimensional life history pattern, while still capturing the essential information. Both Charles Darwin and Alexander von Humboldt used the method of reduced dimensionality to focus ones attention to the key issues while capturing most of the significant information concerning the topic at hand. For example, after reviewing much literature, Darwin concluded that "latitude is a more important element than longitude" for explaining the distribution of organisms (Barrett *et al.* 1987). It was

Humboldt, however, who first used a transect technique to visualize the advantage of reduced dimensionality in explaining observed patterns in distribution (Gayet p. 2284-2287 in Tort 1996). In fisheries science, a classic example of data suitable for reduced dimensionality was presented by Garstang (1909) for the North Sea plaice (*Pleuronectes platessa*, Figure 1 - page 3). Heincke (1913) re-expressed this as a 'law' wherein water depth and/or distance from shore explained most of the observed life history distribution patterns.

The life history characteristics of many species and stocks show generalized two-dimensional patterns, involving water depth and/or distance from shore. For example, FAO (1972) used this approach for many species in their *Atlas of the Living Resources of the Seas*. It is recognized that an inshore/offshore axis may better convey information on structure and processes than an alongshore axis or general geographic map view (Pauly and Lightfoot 1992). A good example of this is demonstrated by comparison of Garstang's map-view of plaice size distribution in the North Sea (Figure 1) with our representation of the same information for the same

Continued on page 4 - Life Histories

The **Sea Around Us** project newsletter is published by the Fisheries Centre at the University of British Columbia. Included with the Fisheries Centre's newsletter *FishBytes*, six issues of this newsletter are published annually. Subscriptions are free of charge.



Our mailing address is: UBC Fisheries Centre, 2204 Main Mall, Vancouver, British Columbia, Canada, V6T 1Z4. Our fax number is (604) 822-8934, and our email address is SeaNotes@fisheries.com. All queries (including reprint requests), subscription requests, and address changes should be addressed to Melanie Power, *Sea Around Us* Newsletter Editor.

The Sea Around Us website may be found at www.fisheries.ubc.ca/Projects/SAUP/index.htm, and contains up-to-date information on the project.




Figure 1: Schematic representation (geographic map view) of the distribution of plaice (*Pleuronectes platessa*) in the North Sea. Mean sizes (cm TL) are given for each depth isobar (modified after Garstang 1909)




Figure 2: Generalised life history pattern by depth zone for plaice in the North Sea (*Pleuronectes platessa*). Depth transect from 53.8°N, 8.6°E to 56.9°N, 3.5°E.

Life Histories - Continued from page 2

species and area (Figure 2 - page 3). Such a transect approach allows the use of icons for key processes, and permits standardization of axis (e.g. log scale), which enables most species or stocks to be directly compared across extensive depth and distance scales. Application of this transect method in the context of the *Sea Around Us Project* will require drawings of similar transects for all important commercial species of the North Atlantic.

The visualization of two-dimensional life history patterns is clearly only a small part in our evaluation of ecosystem effects of fishing (see issue 3, *Sea Around Us* newsletter). Firstly, we will use these transect distributions to help assign catch data to areas such as those described in the classification systems of Large Marine Ecosystems (Sherman and Duda 1999) and 'biogeochemical provinces' (Longhurst 1995). A consensus synthesis approach to these classification systems is being considered by the *Sea Around Us Project* (see Issue 2, Jan/Feb 2000). Secondly, the depth and distance from the coast of major fish population components determines their relative vulnerability to coastal (often small-scale) and offshore (often large-scale) fishing gear and hence potential interactions and conflicts between these different fishery sectors. We will be superimposing the various scales of operation of each fishery sector onto the life history illustrations of each species concerned. Thus, coastal transects of fish distributions will show different species 'connect', through their life history patterns, different

fisheries sectors, such as small with large scale fisheries.

We consider the present approach useful for visualizing the existence, interaction and potential conflicts between different fishery sectors for species or stocks whose life history patterns illustrate the need for improved integration of management of the different fishery sectors. This may apply in particular to rationalization of overcapitalized fisheries. The proposed visualization may be used by management to incorporate the concept of life history interconnectivity between different fishery sectors and may assist in the formulation of more informed policy options for ecosystem-based management of North Atlantic fisheries.

Acknowledgements:

We would like to thank the Environment Project of The Pew Charitable Trusts for their funding of the *Sea Around Us Project*. Thanks also to Drs Tony Pitcher, Reg Watson and Lore Ruttan for assistance with the depth transects and for comments on an earlier version of this article.

References:

Barrett,PH; Gautrey,PJ; Herbert,S; Kohn,D and Smith,S (Editors). 1987. Charles Darwin's Notebooks, 1836-1844: geology, transmutation of species, metaphysical enquiries. British Museum (Natural History), London and Cornell University Press, Ithaca and New York, 747 p.

FAO. 1972. Atlas of the living

resources of the seas. FAO Fisheries Circular No. 126.

Garstang, W. 1909. The distribution of the plaice in the North Sea, Skagerrak and Kattegat, according to size, age, and frequency. Rapp. Council. Int. Explor. Mer. 1:136-138.

Heincke, F. 1913. Investigation on the plaice. General Rapport I. Plaice fishery and protective regulation. Part I. Rapp. P.-v. Reun. Council. Int. Explor. Mer. 17A:1-153.

Longhurst, AR. 1995. Seasonal cycles of pelagic production and consumption. Prog. Oceanog. 36:77-167.

Pauly,D and Lightfoot,C. 1992. A new approach for analyzing and comparing coastal resource systems. NAGA 15(2):7-10.

Sherman,K and Duda,AM. 1999. Large Marine Ecosystems: an emerging paradigm for fishery sustainability. Fisheries 24(12):15-20.

Tort, P. (Editor) 1996. Dictionnaire du Darwinisme et de l' évolution. Volume 2. Presses Univeritaires de France.

Dirk Zeller is a post-doctoral fellow with the Sea Around Us project. Daniel Pauly is Project Leader.